


Juan de Fuca Strait 123°30'W

BACKSCATTER STRENGTH and SHADED SEAFLOOR RELIEF of the

Southern Gulf Islands and San Juan Archipelago, Canada and USA



EDITORS: H. Gary Greene, J. Vaughn Barrie

AUTHORS: Charlie Endris and Kim Picard

ACKNOWLEDGEMENTS: Canadian Hydrographic Service participants R. Hare, K. Czotter, E. Sargent, P. Milner, J. Gagne, C. Lessels, D. Cartwright, K. Lyngberg, and Canadian Coast Guard participants G. Allison and A. Keene

This map was produced by the Center for Habitat Studies at Moss Landing Marine Laboratories in cooperation with Tombolo, the SeaDoc Society, Natural Resources Canada, and the Canadian Hydrographic Service

Digital Cartography by Charlie Endris, Center for Habitat Studies

DESCRIPTIVE NOTES

Backscatter intensity is a measure of sound that is scattered back toward the transmitter by acoustic reflection and scattering, both at the sediment-water interface and within the sediment (volume scattering). Many factors influence the intensity value, among them are: the angle of incidence of the beam, the volume scattered, the seabed slope and the surficial sediment type and roughness. With these factors in mind, backscatter strength datasets are used to determine relative sediment differences within one or many datasets and aid the interpretation of the benthic habitat types. To assure the best interpretation, backscatter images are used in conjunction with other multibeam echosounder bathymetry derivative datasets, such as seafloor shaded relief, slope analysis and bathymetric contours. The multibeam echoesounder bathymetry and backscatter raster datasets, as well as the benthic habitat layer were processed using ESRI□ ArcGIS tools.

The Transboundary region covered by this map series has been divided into four quadrants and this sheet (Sheet 2 of 4; Southern Gulf Islands) covers the Boundary Pass area between Saturna, South Pender and southern Saltspring Islands to the north and northern San Juan, Stuart, Spieden and Waldron islands to the south. Backscatter intensities vary from high (darker areas in mosaic) representing bedrock and coarsegrain sediment (pebbles, cobbles, boulders) to low intensities representing fine-grain sediment (sand and mud). Bedrock is comprised predominantly of hard differentially eroded sedimentary bedrock and medium to low intensities of backscatter represent dynamic bedforms and other sediment deposits in the Boundary Pass, northern Haro Strait, Spieden and President channels area.

